



Claim 57 (Previously Added): The transgenic mammal of claim 56, wherein the matrix metalloproteinase is MMP-13.

Claim 58 (Cancelled)

Claim 59 (Currently Amended): The transgenic mammal of claim ~~58~~ 57, wherein the MMP-13 comprises the sequence of SEQ ID NO: 1 or SEQ ID NO: 21.

Claim 60 (Previously Added): The transgenic mammal of claim 55, wherein the repressor-activator fusion polypeptide is a chimeric tetracycline repressor-VP16 transcription activator polypeptide and the regulatable promoter is a Tn10 sequence linked to a portion of the CMV IE promoter.

Claim 61 (Previously Added): The transgenic mammal of claim 60, wherein the regulatable promoter comprises the sequence of SEQ ID NO: 2.

Claim 62 (Previously Added): The transgenic mammal of claim 55, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 63 (Currently Amended): The transgenic mammal of claim 55, wherein the ~~chondrocyte tissue~~ joint-specific promoter is a Type II collagen promoter.

Claim 64 (Currently Amended): A transgenic rat whose genome comprises:

(a) a nucleotide sequence encoding a constitutively enzymatically active human matrix metalloproteinase that cleaves Type II collagen, wherein the nucleotide sequence encoding the metalloproteinase is operatively linked to a tetracycline-regulatable promoter; and

(b) a nucleotide sequence encoding a repressor-activator fusion polypeptide that binds to the tetracycline regulatable promoter in the absence of tetracycline or a tetracycline analog and does not bind to the regulatable promoter in the presence of tetracycline or tetracycline analog, which nucleotide sequence encoding the repressor-activator fusion polypeptide is operatively linked to a ~~chondrocyte tissue~~ joint-specific promoter,

wherein expression of the metalloproteinase is capable of being repressed in the rat until adulthood, and wherein the metalloproteinase is capable of being expressed in the rat during adulthood to a level sufficient to cause Type II collagen degradation in the joints of the rat.

Claim 65 (Currently Amended): The transgenic rat of claim 64, wherein the matrix metalloproteinase is constitutively enzymatically active MMP-13, the tetracycline-regulatable promoter is tetO7, the repressor-activator fusion polypeptide is tTA, and the ~~chondrocyte tissue~~ joint-specific promoter is a Type II collagen promoter.

Claim 66 (Previously Added): The transgenic rat of claim 64, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup>

degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 67 (Previously Added): A method for producing degradation of Type II collagen in the joints of a transgenic non-human mammal, which method comprises:

(a) maintaining the transgenic mammal of claim 55 in presence of the transcription activator protein-binding compound until adulthood; and

(b) activating expression of the matrix metalloproteinase in the transgenic mammal by withholding the compound from the mammal after the mammal has reached adulthood such that the matrix metalloproteinase degrades Type II collagen in the joints of the transgenic mammal.

Claim 68 (Previously Added): The method according to claim 67, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 69 (Currently Amended): A method for producing degradation of Type II collagen in the joints of a transgenic non-human mammal, which method comprises:

(a) maintaining the transgenic mammal of claim 60 in the presence of tetracycline or a tetracycline analog until adulthood; and

(b) activating expression of the matrix metalloproteinase by withholding the tetracycline or tetracycline analog from the mammal after the mammal has reached adulthood, such that the matrix metalloproteinase degrades Type II collagen in the joints of the transgenic mammal.

Claim 70 (Previously Added): The method according to claim 69, wherein the tetracycline analog is doxycycline.

Claim 71 (Previously Added): The method according to claim 69, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 72 (Previously Added): A method for producing degradation of Type II collagen in the joints of a transgenic rat, which method comprises

(a) maintaining the transgenic rat of claim 64 in the presence of tetracycline or a tetracycline analog until adulthood; and



which nucleotide sequence encoding the transcription activator protein is operatively linked to a ~~chondrocyte-tissue~~ joint-specific promoter;

wherein expression of the metalloproteinase is capable of being repressed in the mammal until adulthood, and wherein the metalloproteinase is capable of being expressed in the mammal during adulthood to a level sufficient to cause Type II collagen degradation in the joints of the mammal.

Claim 76 (Previously Added): The transgenic mammal of claim 75, wherein the matrix metalloproteinase is selected from the group consisting of MMP-1, MMP-8, and MMP-13.

Claim 77 (Previously Added): The transgenic mammal of claim 76, wherein the matrix metalloproteinase is MMP-13.

Claim 78 (Cancelled)

Claim 79 (Currently Amended): The transgenic mammal of claim ~~78~~ 77, wherein the MMP-13 comprises the sequence of SEQ ID NO: 1 or SEQ ID NO: 21.

Claim 80 (Currently Amended): The transgenic mammal of claim 75, wherein the ~~chondrocyte-tissue~~ joint-specific promoter is a Type II collagen promoter.

Claim 81 (Currently Amended): The transgenic mammal of claim 75, wherein the transcription activator protein is a chimeric polypeptide comprising a transactivator domain linked to an ecdysone receptor ligand-binding domain, and wherein the transgenic mammal

further comprises a nucleotide sequence encoding a retinoid X receptor (RXR), which nucleotide sequence encoding RXR is operatively linked to a ~~chondrocyte tissue~~ joint-specific promoter.

Claim 82 (Previously Added): The transgenic mammal of claim 75, wherein the transcription activator protein is a chimeric polypeptide comprising a transactivator domain linked to a progesterone receptor ligand-binding domain.

Claim 83 (Previously Added): The transgenic mammal of claim 75, wherein the transcription activator protein is a chimeric polypeptide comprising a transactivator domain linked to a steroid binding domain.

Claim 84 (Previously Added): The transgenic mammal of claim 75, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 85 (Previously Added): A method for producing degradation of Type II collagen in the joints of a transgenic non-human mammal, which method comprises:

(a) maintaining the transgenic mammal of claim 75 in the absence of the transcription activator protein-binding compound until adulthood; and



(b) activating expression of the matrix metalloproteinase in the transgenic mammal by administering the compound to the mammal after the mammal has reached adulthood such that the matrix metalloproteinase degrades Type II collagen in the joints of the mammal.

Claim 86 (Previously Added): A method for producing degradation of Type II collagen in the joints of a transgenic non-human mammal, which method comprises:

(a) maintaining the transgenic mammal of claim 81 in the absence of ecdysone, an ecdysone analog, or dexamethasone until adulthood; and

(b) activating expression of the matrix metalloproteinase in the transgenic mammal by administering ecdysone, an ecdysone analog, or dexamethasone to the mammal after the mammal has reached adulthood such that the matrix metalloproteinase degrades Type II collagen in the joints of the mammal.

Claim 87 (Previously Added): A method for producing degradation of Type II collagen in the joints of a transgenic non-human mammal, which method comprises:

(a) maintaining the transgenic mammal of claim 82 in the absence of mifepristone (RU 486) until adulthood; and

(b) activating expression of the matrix metalloproteinase in the transgenic mammal by administering mifepristone (RU 486) to the mammal after the mammal has reached

adulthood such that the matrix metalloproteinase degrades Type II collagen in the joints of the mammal.

Claim 88 (Previously Added): The method according to claim 86, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 89 (Previously Added): The method according to claim 87, wherein the Type II collagen degradation results in a loss of proteoglycan, cleavage of type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, or combinations thereof.

Claim 90 (Currently Amended): A method for evaluating the potential of a ~~compound~~ composition to counteract degradation of Type II collagen in joints of a non-human transgenic mammal, which degradation results in a phenotypic change selected from the group consisting of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations thereof, which method comprises:

(a) administering the ~~compound~~ composition to the transgenic mammal of claim 55 in which a phenotypic change has been produced by activation of expression of the metalloproteinase ~~has been activated~~ during adulthood of the transgenic mammal; and

(b) ~~comparing the extent of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change in the mammal to which the ~~compound~~ composition was administered ~~relative to~~ with that of a control transgenic mammal in which the composition was not administered but expression of the metalloproteinase was activated at the same age as it was activated in the animal in which the composition was administered ~~without administering the compound~~,

wherein any less extensive development in the nature or extent of the phenotypic change or any increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change to develop in the mammal that has been administered the ~~compound~~ composition relative to the control mammal, indicates the potential

of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of Type II collagen in joints of a mammal.~~

Claim 91 (Currently Amended): A method for evaluating the potential of a ~~compound~~ composition to counteract degradation of Type II collagen in joints of a non-human transgenic mammal, which degradation results in a phenotypic change selected from the group consisting of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations thereof, which method comprises:

- (a) administering the ~~compound~~ composition to the transgenic mammal of claim 60 in which a phenotypic change has been produced by activation of expression of the metalloproteinase ~~has been activated during adulthood of the transgenic mammal;~~ and
- (b) ~~comparing the extent of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change in the mammal to which the ~~compound~~ composition was administered ~~relative to~~ with that of a control transgenic mammal in which the

composition was not administered but expression of the metalloproteinase was activated at the same age as it was activated in the animal in which the composition was administered ~~without administering the compound,~~

wherein any less extensive development in the nature or extent of the phenotypic change or any increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change to develop in the mammal that has been administered the ~~compound~~ composition relative to the control mammal, indicates the potential of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of Type II collagen in joints of a mammal.~~

Claim 92 (Currently Amended): A method for evaluating the potential of a ~~compound composition~~ to counteract degradation of Type II collagen in joints of a transgenic mouse or rat, which degradation results in a phenotypic change selected from the group consisting of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations thereof, which method comprises:

(a) administering the ~~compound~~ composition to the transgenic ~~mammal~~ rat of claim 64 in which a phenotypic change has been produced by activation of expression of the metalloproteinase ~~has been activated~~ during adulthood of the transgenic rat; and

(b) comparing the extent of ~~loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change in the ~~mammal~~ rat to which the ~~compound~~ composition was administered ~~relative to~~ with that of a control transgenic rat ~~mammal~~ in which the composition was not administered but expression of the metalloproteinase was activated at the same age as it was activated in the animal in which the composition was administered ~~without administering the compound~~,

wherein any less extensive development in the nature or extent of the phenotypic change or any increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change to develop in the ~~mammal~~ rat that has been administered the ~~compound~~ composition relative to the control ~~mammal~~ rat, indicates the

potential of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of~~  
Type II collagen in joints of a mammal.

Claim 93 (Currently Amended): A method for evaluating the potential of a ~~compound~~  
composition to counteract degradation of Type II collagen in joints of a non-human transgenic  
mammal, which degradation results in a phenotypic change selected from the group consisting  
of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change  
in joint function, joint space narrowing, destruction of cartilage, a change in growth plate  
morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations  
thereof, which method comprises:

- (a) administering the ~~compound~~ composition to the transgenic mammal of  
claim 75 in which a phenotypic change has been produced by activation of  
expression of the metalloproteinase ~~has been activated~~ during adulthood of the  
transgenic mammal; and
- (b) ~~comparing the extent of loss of proteoglycan, cleavage of Type II~~  
~~collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space~~  
~~narrowing, destruction of cartilage, a change in growth plate morphology,~~  
~~fibrillation and loss of articular cartilage, or osteophyte formation~~ the  
phenotypic change in the mammal to which the ~~compound~~ composition was  
administered ~~relative to~~ with that of a control transgenic mammal in which the

composition was not administered but expression of the metalloproteinase was  
activated at the same age as it was activated in the animal in which the  
composition was administered without administering the compound,  
wherein any less extensive development in the nature or extent of the phenotypic change or any  
increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen~~  
~~into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction~~  
~~of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or~~  
~~osteophyte formation~~ the phenotypic change to develop in the mammal that has been  
administered the ~~compound~~ composition relative to the control mammal, indicates the potential  
of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of Type II~~  
~~collagen in joints of a mammal.~~

Claim 94 (Currently Amended): A method for evaluating the potential of a ~~compound~~  
composition to counteract degradation of Type II collagen in joints of a non-human transgenic  
mammal, which degradation results in a phenotypic change selected from the group consisting  
of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change  
in joint function, joint space narrowing, destruction of cartilage, a change in growth plate  
morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations  
thereof, which method comprises:





of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of Type II collagen in joints of a mammal.~~

Claim 95 (Currently Amended): A method for evaluating the potential of a ~~compound~~ composition to counteract degradation of Type II collagen in joints of a non-human transgenic mammal, which degradation results in a phenotypic change selected from the group consisting of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations thereof, which method comprises:

- (a) administering the ~~compound~~ composition to the transgenic mammal of claim 82 in which a phenotypic change has been produced by activation of expression of the metalloproteinase ~~has been activated during adulthood of the transgenic mammal;~~ and
- (b) ~~comparing the extent of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change in the mammal to which the ~~compound~~ composition was administered ~~relative to~~ with that of a control transgenic mammal in which the

composition was not administered but expression of the metalloproteinase was activated at the same age as it was activated in the animal in which the composition was administered ~~without administering the compound,~~

wherein any less extensive development in the nature or extent of the phenotypic change or any increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change to develop in the mammal that has been administered the ~~compound~~ composition relative to the control mammal, indicates the potential of the ~~compound~~ composition to counteract the phenotypic change ~~degradation of Type II collagen in joints of a mammal.~~

Claim 96 (Currently Amended): A method for evaluating the potential of a ~~compound~~ composition to counteract degradation of Type II collagen in joints of a non-human transgenic mammal, which degradation results in a phenotypic change selected from the group consisting of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, osteophyte formation, and combinations thereof, which method comprises:

(a) administering the ~~compound~~ composition to the transgenic mammal of claim 83 in which a phenotypic change has been produced by activation of expression of the metalloproteinase ~~has been activated~~ during adulthood of the transgenic mammal; and

(b) ~~comparing the extent of loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change in the mammal to which the ~~compound~~ composition was administered ~~relative to~~ with that of a control transgenic mammal in which the composition was not administered but expression of the metalloproteinase was activated at the same age as it was activated in the animal in which the composition was administered ~~without administering the compound~~,

wherein any less extensive development in the nature or extent of the phenotypic change or any increased length of time required for ~~the loss of proteoglycan, cleavage of Type II collagen into a TC<sup>A</sup> degradation product, a change in joint function, joint space narrowing, destruction of cartilage, a change in growth plate morphology, fibrillation and loss of articular cartilage, or osteophyte formation~~ the phenotypic change to develop in the mammal that has been administered the ~~compound~~ composition relative to the control mammal, indicates the potential

of the compound composition to counteract the phenotypic change ~~degradation of Type II~~  
~~collagen in joints of a mammal.~~